

CORRECTION

Open Access



Correction to: Augmented EPR effect post IRFA to enhance the therapeutic efficacy of arsenic loaded ZIF-8 nanoparticles on residual HCC progression

Xuehua Chen^{1,2†}, Yongquan Huang^{1,2†}, Hui Chen^{1,2}, Ziman Chen¹, Jiaxin Chen¹, Hao Wang^{3*}, Dan Li^{2*} and Zhongzhen Su^{1,2*} 

Correction to: *J Nanobiotechnol* 20:34 (2022)

<https://doi.org/10.1186/s12951-021-01161-3>

Following publication of the original article [1], the authors reported that Fig. 4C was incorrect. The

corrected Fig. 4 and the figure caption are given below. The corrections do not affect the results and conclusions. All authors agree to these corrections and apologize for these errors.

Published online: 02 April 2022

The original article can be found online at <https://doi.org/10.1186/s12951-021-01161-3>.

*Correspondence: haowangfz@foxmail.com; lidan25@mails.sysu.edu.cn; suzhzh3@mail.sysu.edu.cn

†Xuehua Chen and Yongquan Huang contributed equally to this work

¹ Department of Ultrasound, Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai 519000, Guangdong, China

² Guangdong Provincial Key Laboratory of Biomedical Imaging and Guangdong Provincial Engineering Research Center of Molecular Imaging, Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai 519000, Guangdong, China

³ Fine Chemical Industry Research Institute, School of Chemistry, Sun Yat-sen University, Guangzhou 510275, Guangdong, China



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 4 As@ZIF-8 NPs inhibited the invasion, migration and EMT of sublethally heated cells in vitro. Wound healing assay of sublethally heated Hep3B (A) and SMMC7721 cells (B) after incubation with free ATO, As@ZIF-8 NPs and ZIF-8 NPs for 24 h. Scale bar, 800 μ m. Representative images of the migration and invasion of sublethally heated Hep3B (C) and SMMC7721 cells (D) after the indicated treatment and the related quantitative analysis chart (E). Scale bar, 1000 μ m. (F) Protein expression of EMT markers N-cadherin, vimentin and E-cadherin in Hep3B and SMMC7721 cells after the indicated treatment. (ns, no statistical difference; *, $P < 0.05$)

Reference

1. Chen X, Huang Y, Chen H, Chen Z, Chen J, Wang H, Li D, Su Z. Augmented EPR effect post IRFA to enhance the therapeutic efficacy of arsenic loaded ZIF-8 nanoparticles on residual HCC progression. J Nanobiotechnol. 2022;20:34. <https://doi.org/10.1186/s12951-021-01161-3>.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.